

The h -Cobordism Theorem

Fang-Rong Zhan (NC State)

Dec, 2023

Morse functions

All the criticals are non-degenerated.

Criticals

$$(df)_x = 0$$

Non-degenerate

$$\det(\text{Hess}f)_x \neq 0$$

Morse functions

All the criticals are non-degenerated.

Facts

Are isolated. The criticals of a Morse function are isolated.

Do exist. Morse functions form an open dense subset of $C^\infty(M, \mathbb{R})$ in C^2 -topology.

Morse functions, locally

Around a critical x ,

$$f = f(x) - (x^1)^2 - (x^2)^2 - \cdots - (x^\lambda)^2 + (x^{\lambda+1})^2 + \cdots + (x^n)^2$$

for some chart (x^1, \dots, x^n) originated at x .

Morse index

λ

Morse #

of criticals

Morse functions, locally

Around a critical x ,

$$f = f(x) - (x^1)^2 - (x^2)^2 - \dots - (x^\lambda)^2 + (x^{\lambda+1})^2 + \dots + (x^n)^2$$

for some chart (x^1, \dots, x^n) originated at x .

Fact

One can read off a cellular structure of the space from indexes of the criticals, namely,

$$\{\text{criticals of a Morse function}\} \xleftrightarrow{1:1} \{\text{cells of a CW structure}\}$$

an λ -ind critical \mapsto a λ -dim cell

Gradient-like vector field X

- $X \cdot f > 0$ away from criticals;
- $X = \nabla f$ around every critical, where the gradient is induced from the Morse chart.
(We have not introduced a metric on the manifold yet.)

Altering X gives a new Morse function.

Cobordism

A $(n+1)$ -dim cobordism is a 5-tuple $(W^{n+1}; V, V'; h, h')$ of smooth oriented compact manifolds W, V, V' and orientation-preserving diffeomorphisms h, h' such that $\partial W \stackrel{\text{diff}}{\approx} V \sqcup -V'$, where the h - and h' -image is understood, also denoted as

$$V \xrightarrow{W} V'.$$

Product cobordism

$$W \stackrel{\text{diff}}{\approx} V \times [0, 1]$$

Cobordism

A $(n+1)$ -dim cobordism is a 5-tuple $(W^{n+1}; V, V'; h, h')$ of smooth oriented compact manifolds W, V, V' and orientation-preserving diffeomorphisms h, h' such that $\partial W \stackrel{\text{diff}}{\approx} V \sqcup -V'$, where the h - and h' -image is understood, also denoted as

$$V \xrightarrow{W} V'.$$

Product cobordism

$$W \stackrel{\text{diff}}{\approx} V \times [0, 1]$$

Question

Under what topological conditions of W , can we identify it as a product cobordism?

Morse theory on a cobordism

Fact

Relative Morse function does exist.

$V \xrightarrow{W} V'$ admits a Morse function f , with $f(0) = V, f(1) = V'$.

Gradient-like vf does exist.

With respect to f , there admits a gradient-like vector field.


Important observation

If there are no criticals of f , then it **is** a product cobordism.

Question updated

Under what topological conditions of W , can we eliminate all the criticals?

Anatomy of criticals



Anatomy of criticals

Let $g : OD^n \overset{\text{diff}}{\approx} U$ open around p . Let V_0, V_1 be level sets near p with $f(V_0) < f(p) < f(V_1)$.

characteristic embedding

$$\varphi_L : S^{\lambda-1} \times OD^{n-\lambda} \rightarrow V_0 : (u, \theta v) \mapsto g(u \cosh \theta, v \sinh \theta)$$

$$\varphi_R : OD^{\lambda-1} \times S^{n-\lambda} \rightarrow V_1 : (\theta u, v) \mapsto g(v \sinh \theta, u \cosh \theta)$$

left-hand sphere S_L / right-hand sphere S_R

$$\varphi_L(S^{\lambda-1} \times 0), \varphi_R(0 \times S^{n-\lambda-1}).$$

left-hand disk D_L / right-hand disk D_R

The union of integral curves of X ending at p (resp. starting at p).

Surgery of type $(\lambda, n - \lambda)$

We can use surgery to describe the change of characteristic embedding from the left-hand to the right-hand.

$$\chi(V, \varphi) = (V - \varphi(S^{\lambda-1} \times 0)) \sqcup (OD^\lambda \times S^{n-\lambda-1}) / \varphi(u, \theta v) \sim (\theta u, v)$$

And use cobordism to trace this surgery.

Theorem

There exists an elementary cobordism (i.e. Morse $\# = 1$) W such that

$$V \xrightarrow{W} \chi(V, \varphi).$$

Lesson A critical is essentially a cobordism. Can we split them?

Rearrangement of criticals

For criticals p, p' index $\lambda \geq \lambda'$ with $f(p) < f(p')$, there is enough room to alter X to move $S_R(p)$ out of the way of $S_L(p')$ at any level set V_c between $f(p)$ and $f(p')$, since $\dim S_R(p) + \dim S_L(p') = (n - \lambda - 1) + (\lambda' - 1) < n - 1 = \dim V_c$.

Theorem


There exists a Morse function on the cobordism $V \xrightarrow{w} V'$ such that

- $f(V) = -\frac{1}{2}, f(V') = n + \frac{1}{2}$
- $f(p) = \text{index}(p), \forall \text{critical } p$

These functions are called self-indexing, and we can split it as compositions of cobordisms $c_0 \circ \cdots \circ c_\lambda \circ \cdots \circ c_n$, with only λ -index's in c_λ .

Suppose in the following that all Morse functions are self-indexing.

Cancellation of criticals



Cancellation of criticals

Always alter X to make the intersection transverse. We can cancel a pair (p, p') of criticals of index $\lambda, \lambda + 1$, provided that ...

Single-point intersection


If S_R is transversely intersected with S'_L with a single point, then the cobordism is a product cobordism.

Some hard differential topology works are required.

Cancellation of criticals

Always alter X to make the intersection transverse. We can cancel a pair (p, p') of criticals of index $\lambda, \lambda + 1$, provided that ...

Whitney's trick, locally



Globally

Requires $\pi_1 M = 0, \pi_1 V_{a-1}(\mathbb{R}^{a+b-2}) = 0$ ($\Leftarrow \dim M = a + b - 2 \geq 6$).

Eliminating middle index criticals

Let $W_\lambda = c_0 \circ \cdots \circ c_\lambda$, $V = W_{-1} \subset W_0 \subset \cdots \subset W_n$. We can interpret the homology generators as the left-hand disks, namely,

$$H_\lambda(W, V) \cong \bigoplus_{\text{left-hand disks}} \mathbb{Z}\langle D_L^\lambda \rangle \cong H_\lambda(W_\lambda, W_{\lambda-1}),$$

Let $C_\lambda := H_\lambda(W_\lambda, W_{\lambda-1})$ and

$$\partial_\lambda : C_\lambda \rightarrow C_{\lambda-1} : [D_L] \mapsto \sum S'_R \cdot S_L[D_L^{\lambda-1}]$$

is given by the induced differential of long exact sequence of the triad $(W_\lambda, W_{\lambda-1}, W_{\lambda-2})$.

Fact

$$H_*(C_\bullet) \cong H_*(W, V)$$

Eliminating middle index criticals

$$H_\lambda(W, V) \cong \bigoplus_{\text{left-hand disks}} \mathbb{Z}\langle D_L^\lambda \rangle \cong H_\lambda(W_\lambda, W_{\lambda-1}),$$

- If W is a product cobordism, then $H_*(W, V) = 0$.
- Conversely, if $H_*(W, V) = 0$, then $H_*(C_\bullet) = 0$ and therefore C_\bullet is exact, which means every critical is pairing up with another adjacently indexed critical.

Lesson

The criticals' data is essentially the homology data.

Question twice updated

Can $H_*(W, V) = 0, \pi_1 W = 0, \dim W \geq 6$ imply that W is a product cobordism?

Facts that the elimination works generally

- All 0-indexed criticals can be canceled with equal number of 1-indexed's.
- All 1-index's can be traded for equal number of 3-index's, provided $\pi_1 V = 0$.
- Replace f by $-f$ to eliminate 0-coindex's and 1-coindex's.

Yes! from h -cobordism theorem

For $V \xrightarrow{W} V'$, if

- 1) $\pi_1 W = \pi_1 V = 0$;
- 2) h -cobordism;
- 3) $\dim W \geq 6$;

then $W \overset{\text{diff}}{\approx} V \times [0, 1]$.

h -cobordism

V, V' are deformation retract of W .

Remark

Under condition 1),
 $h\text{-cob} \iff H_*(W, V) = 0$.

Characteristic of $D^{\geq 6}$

$\pi_1 W = \pi_1 \partial W = 0$, W compact. TFAE:

1) $W \stackrel{\text{diff}}{\approx} D$

2) $W \stackrel{\text{homeo}}{\approx} D$

3) W retractable

4) $H_* W \cong H_* pt$

1) \implies 2) \implies 3) \implies 4) is obvious. For 4) \implies 1):

Let $D \hookrightarrow W$ be an embedding of a disk, Since

$$H_*(W - \text{int} D, \partial D) \cong H_*(W, D) = 0,$$

by the h -cobordism theorem, $\emptyset \xrightarrow{W - \text{int} D} \partial D$ is a product cobordism. Combining with $\partial D \xrightarrow{D} \emptyset$, we have $W \stackrel{\text{diff}}{\approx} D$.

Generalized Poincaré Conjecture

If $\pi_1 V^n = 0$, $H_* V \cong H_* S^n$, $n \geq 5$, then $V \stackrel{\text{homeo}}{\approx} S^n$.

Let $D^n \hookrightarrow V$ be an embedding of a n -disk.

$$H_*(V - \text{int} D, \emptyset) \cong H^{n-*}(V - \text{int} D, \partial D) \cong H^{n-*}(V, D) \cong \begin{cases} \mathbb{Z}, & * = 0 \\ 0, & * \neq 0 \end{cases} \cong H_* pt$$

By the characteristic of $D^{\geq 6}$, we have $V - \text{int} D \stackrel{\text{diff}}{\approx} D_0^n$, that is $V^n \stackrel{\text{diff}}{\approx} D^n \cup_h D_0^n$, a so-called "twisted sphere", where $h : \partial D \rightarrow \partial D_0$ is the attaching map. Let $g_0 : D_0 \hookrightarrow S^n$ be a

embedding, then $g(u) = \begin{cases} g_0(u), & u \in D_0 \\ \sin \frac{\pi t}{2} g_0(h^{-1}u) + \cos \frac{\pi t}{2} e_{n+1}, & u \in D \end{cases}$ is a homeomorphism.

Thank you for your attention!

Questions?